

## UNITED STATES DEATHS OF PEOPLE WITH "AIDS" DIAGNOSES

|   | Number  | Percent |
|---|---------|---------|
| Cumulative total from beginning of epidemic   | 583,298 | 100.0   |
| <b>YEAR</b>                                   |         |         |
| From before 1981 through 1986<br>(Before AZT) | 24,717  | 4.2     |
| From 1987 through 2007<br>(After AZT)         | 558,581 | 95.8    |

This table is constructed from data in two CDC HIV/AIDS Surveillance Reports: the year-end report through December 1997 and the year-end report through 2007. The former gave deaths by year for 1981-1997. The latter gave cumulative totals through 2007. (About a decade ago the CDC stopped reporting "AIDS" deaths by year, including past years, which made it necessary to combine these two reports.)

Observe that 96% of the "AIDS deaths" occurred after AZT was introduced in 1987 the year that "AIDS" incidence peaked and the CDC greatly expanded the surveillance definition.

What did all these people die of? **Poisons** (both "recreational" and pharmaceutical) and lethal diagnoses and counselling. They were prescribed toxic and worthless drugs that they didn't need. They were never told what the real risk factors are for AIDS-indicator diseases. (Heroin users were never told to stop injecting heroin, only to use clean needles. Gay men were told it was safe to use poppers.) They were all told that they would die from an invariably fatal disease.

Note that most deaths among "HIV-positive" individuals (according to the unvalidated HIV-antibody tests) now occur before they have acquired an "AIDS" diagnosis. These are called "death before diagnosis". For years in Massachusetts the leading cause of death among patients with an "HIV-positive" diagnosis has been liver cancer, caused by the various "anti-viral" drugs. These are not counted as "AIDS" deaths. Therefore, the above data greatly understate the lethality of the "anti-viral" drugs, which are the leading cause of "HIV-positive" deaths, as well as "AIDS" deaths.

John Lauritsen <http://paganpressbooks.com/jpl/AIDS.HTM>

## RELATIVE "AIDS" DEATHS: US & UK

|                            | US          | UK         |
|----------------------------|-------------|------------|
| Est. Population 2008       | 304,059,724 | 58,789,194 |
| "AIDS" Deaths through 2007 | 583,298     | 18,787*    |
| Proportion: Deaths/Pop.    | 0.19%       | 0.03%      |
| Proportions Ratio: US/UK   |             | 6.3        |

\* Includes "deaths before diagnosis" as well as "AIDS" deaths. The American statistics count only those with an "AIDS" diagnosis.

Comment: Taking population differences into account, there have been over six times as many "AIDS" deaths in the United States as in the United Kingdom. Probably the Meditel documentaries and Neville Hodgkinson's *Sunday Times* articles, which warned about the dangers of AZT, are partly responsible for this large disparity. No television documentary critical of AIDS orthodoxy has ever aired in the US. Nor has any major US newspaper ever fairly covered AIDS criticism.

John Lauritsen <http://paganpressbooks.com/AIDS.HTM>