

Study Group AIDS-therapy c/o Felix de Fries Juliastr. 8 8032 Zürich felix.defries@gmail.com

To people affected
Their doctors and carers
To Media

Zürich, 12th August 2020

ref. **Vaccines or antiviral substances**

Dear Sir/Madam

Since the beginning of the Covid-19 pandemic, we are taught that only a vaccine, which should be available by the end of this year could resolve the problem. Such a vaccine could be inactivated, attenuated, protein-based, RNA-based, DNA-based, virus-like particle based or viral vector based, as shown by 363 patent applications. In addition to the search for a vaccine, antiviral substances were also sought from the start including already existing ones, that have been developed against viral infections such as Ebeola, MERS and influenza, broad spectrum antivirals, entry inhibitors, therapeutic antibodies, cytokines, mRNA therapeutics to unbind expression and transcription and micro particles, that should block viral proteins. After it is foreseeable that an effective vaccine against a constantly mutating RNA-virus, which can exist in one person in the form of different strains and mutants, cannot be found on the short term, even a company such as ROCHE, who has made the test to measure an active viral infection available world-wide apparently doubts the making of an effective vaccine in this time and wants to participate in the search for a new antiviral substance by a newly built research team.

[Research and development on therapeutic agents and vaccines for COVID-19 and related human coronavirus diseases](#)

[HTML\] acs.orgFull View](#)

Preventive therapies against the severe course of Covid-19, which takes place due to pre-existing illness such as hypertension, obstructive pulmonary diseases, cardio-vascular diseases with thrombosis risk and diabetes, all of which occur under high air pollution are not an issue for Big Pharma and for the health care institutions. They do not disclose with their daily information on new infections, how many of those tested positive for SARS Cov-2 are going through a difficult course. (New statistical data from Switzerland now reveals that only 10% of positive tested need hospitalization.) The fact, that two thirds of them suffer from malnutrition, as it has been demonstrated in studies from Italy is furthermore no reason for according preventive treatment in vulnerable groups.

[\[HTML\] Prevalence of malnutrition and analysis of related factors in elderly patients with COVID-19 in Wuhan, China](#)

[\[HTML\] nature.com](#)

[\[HTML\] Elderly at time of COroNaVirus disease 2019 \(COVID-19\): possible role of immunosenescence and malnutrition](#)

[\[HTML\] springer.com](#)

[Enhancing immunity in viral infections, with special emphasis on COVID-19: A review](#)

[\[HTML\] nih.gov](#)

[Early nutritional supplementation in non-critically ill patients hospitalized for the 2019 novel coronavirus disease \(COVID-19\): Rationale and feasibility of a shared ...](#)

[\[HTML\] nih.gov](#)

[\[HTML\] Clinical significance of nutritional risk screening for older adult patients with COVID-19](#)

[\[HTML\] nature.com](#)

[Clinical characteristics of 25 death cases with COVID-19: a retrospective review of medical records in a single medical center, Wuhan, China](#)

[\[PDF\] sciencedirect.com](#)

[COVID-19: The Inflammation Link and the Role of Nutrition in Potential Mitigation](#)

[\[PDF\] mdpi.com](#)

[Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis](#)

[HTML\] ijidonline.com](#)

[Influenza and obesity: its odd relationship and the lessons for COVID-19 pandemic](#)

[\[HTML\] nih.gov](#)

For people living in countries of the developing world, who have to earn a daily income for their meals by informal work the declared distance keeping rules represent an unattainable challenge that requires answers on various levels.

[Multidimensional poverty and COVID-19 risk factors: A rapid overview of interlinked deprivations across 5.7 billion people](#)

[\[PDF\] ox.ac.uk](#)

[\[PDF\] Slum health: arresting COVID-19 and improving well-being in urban informal settlements](#)

[\[PDF\] springer.com](#)

State health system officials continue to pretend in their epidemic information policies, that Covid-19 is all about avoiding contagion which according to their statements can cause serious fatal disease in everyone.

After becoming increasingly visible that in today's work- and consumer societies, the spread of a micro-droplet infection cannot be completely prevented by social distancing, contact tracing and quarantine, statal institutions such as Swissmedic, without a clear proof of effect and of adverse effects approve the administration of nucleoside-analogue antivirals such as Remdesivir, said to shorten the stay in intensive care units, even as it cannot prevent fatal outcomes in severely ill persons.

The previously agreed rules on clinical trials with ill patients have been quickly thrown overboard.

https://www.swissmedic.ch/swissmedic/de/home/news/coronavirus-covid-19/erweitert_einsatz_remdesivir.html

That non-patented, natural substances such as Azetylcysteine can prevent complications and significantly shorten the stay on intensive care units does not seem to be an issue in this context.

[\[HTML\] Safety and efficacy of N-acetyl-cysteine for prophylaxis of ventilator-associated pneumonia: a randomized, double blind, placebo-controlled clinical trial](#)

Like at the beginning of HIV-AIDS, when the representatives of pharma-medicine and their political lobbies suppressed studies on preventive immune-system bolstering therapies (including those on Azetyl-Cystein) and pushed through the administration of nucleoside analogue substances such as AZT, known to be seriously harmful, with the effect that thousands (first mainly homosexual males) died from this antiviral monotherapy before 10 years later a combi therapy was launched with reduced doses of these substances and 25 years later the Pre-Exposition Therapy (PreP), able to suppress a positive test-result in persons suffering of syphilis, Chlamydia and Gonorrhoea infections were administrated. Such as with the HIV-AIDS-paradigm, by which the severe course of more than 30 mostly endemic diseases (from Tuberculosis, over Hepatitis A,B and C, Malaria, Herpes I and II, HPV, Syphilis,

Gonorrhoea and Chlamydia) were traced back to the HIV-retrovirus, told to cause a fatal infection, now a new scenario of fear should allow again a billion selling business with antiviral substances, vaccines and tests.

Mitochondrial toxicity of antiviral drugs

<https://www.nature.com/articles/nm0595-417>

FDA Documents Show Fraud In AZT Trials

<https://paganpressbooks.com/jpl/FRAUD.PDF>

By means of the obligatory carrying of masks allegedly everyone should get a musszle which makes the face non-recognizable. Anyone not having a steady partner now cannot experience closeness. By means of these measures, we are supposed to be ready for the intake of immature antivirals and vaccines to be without its proven intake no longer allowed to attend events or to work.

Felix de Fries

[AIDS at viral load zero](#)

[AIDS at viral load zero studies and links](#)